
Evaluation of a Prolonged Prothrombin Time

Joshua L. Hood and Charles S. Eby*

Department of Pathology and Immunology, Washington University School of Medicine, St. Louis, MO.

*Address correspondence to this author at: Department of Pathology and Immunology, Washington University School of Medicine, St. Louis, MO 63110. E-mail eby@wustl.edu.

CASE DESCRIPTION

A 47-year-old African American woman was evaluated for a prolonged prothrombin time (PT) result obtained before she underwent right total hip arthroplasty. The patient had no history of gastrointestinal or intracranial bleeding, epistaxis, or hemarthrosis. However, she reported a tendency toward easy limb bruising and menorrhagia, which required iron supplementation. She had a negative family history of abnormal bleeding. Initial laboratory studies included findings within reference intervals for complete blood cell count and activated partial thromboplastin time (aPTT) (30.8 s, reference interval 23–36 s), prolonged PT (20.3 s, reference interval 11.0–15.0 s), and International Normalized Ratio (INR) (1.78, reference interval 0.9–1.2). No preanalytical artifacts were identified, and the result of a repeat PT was also prolonged.

Questions to Consider
<ul style="list-style-type: none">• What are some of the preanalytical causes of inaccurate PT and aPTT results?
<ul style="list-style-type: none">• If the prolonged PT is an accurate result, what coagulation tests should be done to determine the specific cause?
<ul style="list-style-type: none">• If these results are due to an inherited coagulation deficiency, which clotting factor is most likely to be involved?

Final Publication and Comments

The final published version with discussion and comments from the experts will appear in the April 2008 issue of *Clinical Chemistry* in approximately 3–4 weeks. To view the case and comments online, go to <http://www.clinchem.org/content/vol54/issue4/> and follow the link to the Clinical Case Study and Commentaries.

Educational Centers

If you are associated with an educational center and would like to receive the cases and questions 3-4 weeks in advance of publication, please email clinchem@aacc.org.

AACC is pleased to allow free reproduction and distribution of this Clinical Case Study for personal or classroom discussion use. When photocopying, please make sure the DOI and copyright notice appear on each copy.

The Clinical Case Study for April can be found at
<http://www.clinchem.org/CCS/casestudies.shtml>.

All previous Clinical Cases Studies can also be accessed and downloaded online at or
<http://www.aacc.org/AACC/resources/casestudies/>.

AACC is a leading professional society dedicated to improving healthcare through laboratory medicine. Its nearly 10,000 members are clinical laboratory professionals, physicians, research scientists, and others involved in developing tests and directing laboratory operations. AACC brings this community together with programs that advance knowledge, expertise, and innovation. AACC is best known for the respected scientific journal, *Clinical Chemistry*, the award-winning patient-centered web site *Lab Tests Online*, and the world's largest conference on laboratory medicine and technology. Through these and other programs, AACC advances laboratory medicine and the quality of patient care.